

## AF 594 tyramide



<http://cn.lumiprobe.com/p/af-594-tyramide-6>

Thyramide signal amplification (TSA) is the most versatile and effective way to enhance the intensity of the fluorescent signal, used in immunohistochemistry (IHC), immunocytochemistry (ICC), and fluorescence *in situ* hybridization (FISH). The TSA method is based on the ability of horseradish peroxidase (HRP) in the presence of low concentrations of hydrogen peroxide to convert a labeled tyramine-containing substrate into an oxidized, highly reactive free radical that covalently binds to the tyrosine residues of protein molecules located next to it.

Compared to conventional procedures, the TSA method increases the sensitivity of immunofluorescent detection of target molecules by more than 100 times, making it particularly suitable for detecting low-concentration targets. In applications where increased detection sensitivity is not required, TSA can significantly reduce antibody or probe concentrations without loss of signal intensity, thereby reducing background staining due to cross-reactivity or non-specific binding of antibodies.

Since the binding of the tyramide label is covalent, tyramides of different dyes can be used in several sequential rounds of TSA staining to detect multiple targets in the same sample.

This tyramide is a conjugate of the water-soluble red fluorescent dye AF 594. AF 594 tyramide is a component of many tyramide signal amplification (TSA) kits. It can be used with any antibody or other molecules (streptavidin, etc.) conjugated to HRP to stain cells and tissues by immunofluorescence methods.



外观:

分子 943.15

量:

分子 C<sub>49</sub>H<sub>59</sub>N<sub>4</sub>O<sub>11</sub>S<sub>2</sub>

式:

溶解

度:

质量

控制:

储存

条件:

法律 本产品仅供研究目的提供和销售。 本产品并未经过食品、药品、医疗器械、化妆品等领域的安全性和效力测试, 且未经明示或暗示授权用于其他任何用途, 包括但不限于体外诊断、人类或动物用途, 以及商业用途。

激发/ 586

吸收

极大

值,

纳米:

ε, 摩 105000

尔吸

光系

数 **art**

发射 613  
极大  
值,  
纳米:  
荧光 0.77  
量子  
产率:  
CF<sub>260</sub>: 0.28  
CF<sub>280</sub>: 0.51