


## Alkyne-PEG4-maleimide

<http://cn.lumiprobe.com/p/alkyne-peg4-maleimide>

When proteins or other sensitive biomolecules need to be conjugated, polyethyleneglycol (PEG) linkers are a great choice. They provide hydrophilicity and maintain separation between conjugated molecules. This linker based on PEG4, has length of 14Å (1.4 nm). Heterobifunctional linkers carrying different reactive groups can be used to prepare cross-conjugates between different molecules.

Maleimide functional group is especially useful, because they can label proteins site specifically. Most proteins have only limited number of cysteine residues that selectively react with maleimides. Alkyne group can be then conjugated with azides in a selective copper-catalyzed process, CuAAC (copper-catalyzed alkyne-azide cycloaddition).



外观:  
质 311.3  
谱  
M+  
增  
量:  
分 311.33  
子  
量:  
CAS 1262681-30-0  
编  
号:  
分 C<sub>15</sub>H<sub>21</sub>NO<sub>6</sub>  
子  
式:  
溶  
解  
度:  
质  
量  
控  
制:  
储  
存  
条  
件:  
法 本产品仅供研究目的提供和销售。 本产品并未经过食品、药品、医疗器械、化妆品等领域的安全性和效力测试, 且未经明示或暗示授权用于其他任何用途, 包括但不限于体外诊  
律 断、人类或动物用途, 以及商业用途。  
声  
明: