


Biotin-X-NHS ester

<http://cn.lumiprobe.com/p/biotin-x-nhs-ester>

Biotin may be conjugated to many proteins while maintaining the biological activity. The biotinylated probe is usually detected by avidin, streptavidin due to their effective binding which is used in various applications such as affinity chromatography, ELISA and Western Blot, fluorescence-activated cell sorting (FACS), intracellular labeling.

Biotin-X-NHS ester is a derivative with C6-spacer to reduce steric effect when binding amino acids, peptides, or proteins by reacting with primary amines. This compound is used to attach biotin to primary amines under alkaline conditions (pH 8-9).



外观:

质谱 339.2

M+

增量:

分子量: 454.55

量:

CAS 72040-63-2

编号:

分子式: C₂₀H₃₀N₄O₆S

式:

IUPAC (2,5-dioxopyrrolidin-1-yl) 6-[5-[(3aS,4S,6aR)-2-oxo-1,3,3a,4,6,6a-hexahydrothieno[3,4-d]imidazol-4-yl]pentanoylamino]hexanoate

名称:

溶解

度:

质量

控制:

储存

条件:

法律 本产品仅供研究目的提供和销售。 本产品并未经过食品、药品、医疗器械、化妆品等领域的安全性和效力测试，且未经明示或暗示授权用于其他任何用途，包括但不限于体外诊断、人类或动物用途，以及商业用途。