


## C26:0-lysophosphatidylcholine (LPC)

<http://cn.lumiprobe.com/p/c26-lpc-1213783-80-2>

Lysopc(26:0), also known as LysoPC a C26:0, is classified as a member of the 1-acyl-sn-glycero-3-phosphocholines. 1-acyl-sn-glycero-3-phosphocholines are glycerophosphocholines in which the glycerol is esterified with a fatty acid at O-1 position, and linked at position 3 to a phosphocholine. Lysopc(26:0) can be used as internal standard LC-MS studies.

Lysophosphatidylcholines (LPCs) specifically bind to G protein-coupled receptors GPR119, GPR40, GPR55 and GPR4. Binding of LPCs to GPR119, GPR40 and GPR55 induces intracellular calcium mobilization and leads to increased glucose-stimulated insulin secretion in different cell systems. In blood or plasma LPCs are bound mainly to albumin and to a lesser extent to lipoproteins.

Inflammation, cell damage and other pathophysiological conditions can profoundly alter the ratio of free to albumin bound LPC through increased production of LPC or decreased plasma levels of albumin. In particular, lower levels of albumin (hypoalbuminemia) lead to lower levels of LPC in the blood. Lysopc(26:0) is a biomarker for X-linked Adrenoleukodystrophy (X-ALD). Lysopc(26:0) could be used as auxiliary marker for diagnosis of peroxisomal  $\beta$ -oxidation disorders.



外观: 白色粉末

分子 635.90

量:

CAS 1213783-80-2

编号:

分子  $C_{34}H_{70}NO_7P$

式:

IUPAC [(2R)-3-hexacosanoyloxy-2-hydroxypropyl] 2-(trimethylazaniumyl)ethyl phosphate

名称:

质量

控制:

储存

条件:

法律 本产品仅供研究目的提供和销售。 本产品并未经过食品、药品、医疗器械、化妆品等领域的安全性和效力测试，且未经明示或暗示授权用于其他任何用途，包括但不限于体外诊断、人类或动物用途，以及商业用途。