

## Exo-BCN-PNP

<http://cn.lumiprobe.com/p/exo-bcn-pnp-carbonate>


BCN-PNP carbonate is a tool to introduce BCN moiety via carbamate linkage with primary amino groups of substrate. Reaction conditions is similar to NHS-amino linkage but PNP ester provides less off-target hydrolysis and higher conjugation yield forming hydrolytically stable bond with substrate.

Bicyclononyne (BCN) is one of the most reactive cyclooctynes for copper-free click chemistry. Both exo- and endo- isomers are active and have near the same power of reactions konstants [1,2]. BCN reacts even faster than DBCO [1] with aromatic azides and comparing to dibenzofused systems has two advantages. First, it contains a plane of symmetry which prevents the formation of mixtures of stereoisomeric products. Second, BCN provides lower lipophilicity which is typically more beneficial when reaction is performed in aqueous solutions. Unlike DBCO, BCN reactivity is not only limited to azides (SPAAC) and nitrones (SPANC), but also covers [tetrazines](#) (IEDDA) [2] and more recently tetrazoles (photoclick) [3] providing exceptional reaction rates.

[1] Jan Dommerholt et al. Readily Accessible Bicyclononynes for Bioorthogonal Labeling and Three-Dimensional Imaging of Living Cells. *Angewandte Chemie*. 2010. 49(49). 9422-9425.

[2] Wagner et al. Origin of Orthogonality of Strain-Promoted Click Reactions. *Chemistry*. 2015. 21(35).12431-12435.

[3] Gangam Srikanth Kumar et al. Superfast Tetrazole-BCN Cycloaddition Reaction for Bioorthogonal Protein Labeling on Live Cells. *Journal of the American Chemical Society*. 2022. 144(1). 57-62.



外观

分子量 315.32

分子式

$C_{17}H_{17}NO_5$

溶解度

质

量 NMR  $^1H$  和 HPLC-MS (95+%)

控制

储存 收到后 -20°C 避光保存 24 个月。运输: 室温下最多可保存3周。干燥。

条件

法律声明 本产品仅供研究目的提供和销售。本产品并未经过食品、药品、医疗器械、化妆品等领域的安全性和效力测试, 且未经明示或暗示授权用于其他任何用途, 包括但不限于体外诊断、人类或动物用途, 以及商业用途。