

Fluo-4 AM, green fluorescent calcium indicator

<http://cn.lumiprobe.com/p/fluo-4-am>



Fluo-4 AM is a cell-permeable Ca^{2+} -indicator that is metabolized by intracellular esterase, leading to a bright green fluorescent signal upon Ca^{2+} -binding (excitation/emission λ at 494/506 nm). Fluo-4 AM is used for visualization and measurement of intracellular Ca^{2+} . It is well suited for fluorometric and imaging applications such as microscopy, flow cytometry, spectrofluorometry, and fluorometric high-throughput microplate screening assays [1].

Fluo-4 AM is similar in structure and spectral properties to the widely used Ca^{2+} -indicator, Fluo-3, but it has certain advantages over Fluo-3, such as brighter fluorescence emission, high rate of cell permeation, and a K_d for Ca^{2+} in buffer of 345 nM. Because of its higher fluorescence emission intensity, Fluo-4 AM can be used at lower intracellular concentrations, making its use less toxic for live cells.

As Fluo-4 AM does not covalently bind to cellular components, it may be actively effluxed from the cell by organic anion transporters. *In vivo* cell imaging with Fluo-4 AM is usually performed within one or two hours after loading, but the dye can be re-loaded to cells if it is needed. Fluo-4 AM can also be fixed *in situ* by [EDC/EDAC](#) for downstream immunofluorescence studies.

Fluo-4 AM has low solubility in the water. It is recommended to prepare 1 mM stock solution in [labeling grade DMSO](#) prior to cell loading. Use the final concentration of 1-5 μM and incubation at 37 °C for 15-60 min as a start point of your protocol.

[1] Gee K.R. et al. Chemical and physiological characterization of fluo-4 Ca^{2+} -indicator dyes. *Cell Calcium*. 2000. 27(2). 97-106.



外观:

分子 1096.95

量:

CAS 273221-67-3

编号:

分子 $\text{C}_{31}\text{H}_{50}\text{F}_2\text{N}_2\text{O}_{23}$

式:

IUPAC N-[4-[6-[(Acetyloxy)methoxy]-2,7-difluoro-3-oxo-3H-xanthen-9-yl]-2-[2-[2-bis[2-[(acetyloxy)methoxy]-2-oxoethyl]amino]-5-methylphenoxy]ethoxy]phenyl]-N-[2-[(acetyloxy)methoxy]-2-oxoethyl]glycine
名称: (acetyloxy)methyl ester

溶解

度:

质量

控制:

储存

条件:

激发/ 494

吸收

极大

值,

纳米:

