

FluoriCa-8 AM, green fluorescent calcium indicator

<http://cn.lumiprobe.com/p/fluo-8-am>



FluoriCa-8 AM is a cell-permeable Ca^{2+} -indicator that is metabolized by intracellular esterase, leading to a bright green fluorescent signal upon Ca^{2+} -binding (excitation/emission λ at 490/514 nm). FluoriCa-8 AM is used for visualization and quantifying intracellular Ca^{2+} . It is well suited for fluorometric and imaging applications such as microscopy, flow cytometry, spectrofluorometry, and fluorometric high-throughput microplate screening assays.

FluoriCa-8 AM is similar in structure and spectral properties to the Ca^{2+} indicators Fluo-3 AM and Fluo-4 AM but has the brightest fluorescence compared to them (two times brighter than Fluo-4 and four times brighter than Fluo-3). The K_d of FluoriCa-8 AM for Ca^{2+} is about 389 nM. With its highest fluorescence intensity, FluoriCa-8 AM is ideal for applications where the concentration of dye loaded into cells is required to be minimized. Unlike Fluo-3 AM and Fluo-4 AM, which require cells to be incubated at 37 °C, FluoriCa-8 AM can be loaded into cells at room temperature.

As FluoriCa-8 AM does not covalently bind to cellular components, it may be actively effluxed from the cell by organic anion transporters. *In vivo* cell imaging with FluoriCa-8 AM is usually performed within one or two hours after loading, but the dye can be re-loaded to cells if it is needed. To determine the cellular localization of the dye, FluoriCa-8 AM can be fixed *in situ* with [EDC/EDAC](#) and counterstained immunochemically using anti-BAPTA antibodies.

FluoriCa-8 AM has low solubility in the water. It is recommended to prepare 1 mM stock solution in [labeling grade DMSO](#) prior to cell loading. Use the final concentration of 1-5 μM and incubation at RT for 15-60 min as a start point of your protocol.

FluoriCa-8 AM solutions in DMSO may be stored at -20 °C for up to 2 months.



外观:
分子 1060.97
量:
CAS 1345980-40-6
编号:
分子 $\text{C}_{51}\text{H}_{52}\text{N}_2\text{O}_{23}$
式:
溶解度:
质量控制:
储存条件:

法律声明: 本产品仅供研究目的提供和销售。 本产品并未经过食品、药品、医疗器械、化妆品等领域的安全性和效力测试, 且未经明示或暗示授权用于其他任何用途, 包括但不限于体外诊断、人类或动物用途, 以及商业用途。

激发 494

吸收
极大
值，
纳
米：

发射 517

极大
值，
纳
米：