

H2DCFDA (2',7'-dichlorodihydrofluorescein diacetate)

<http://cn.lumiprobe.com/p/h2dcfda>



H₂DCFDA (2',7'-dichlorodihydrofluorescein diacetate) is a common reagent used for investigating the production of reactive oxygen species in living cells.

H₂DCFDA is a non-fluorescent fluorescein derivative (its reduced acetylated form). The reagent begins to emit fluorescence only after cleavage of acetyl groups and the reagent's oxidation in the cell while converting to 2',7'-dichlorofluorescein. This is a bright green-fluorescent dye (absorption maximum 511 nm, fluorescence maximum 533 nm). This reagent can be used for assays in living cells and is not compatible with sample fixation.

Acetyl groups in H₂DCFDA increase its lipophilicity and improve its cell membrane permeability. Once got into the cell, the dye is deacetylated by cell esterases, thus becoming charged and better fixed inside the cell. Oxidation with reactive oxygen forms results in the formation of a fluorescent product (2',7'-dichlorofluorescein) and can be detected using various methods, for example with a flow cytometer, plate reader, or fluorescent microscope.

Recommendations for using the reagent:

- Use a freshly prepared solution of the reagent (the working solution is not intended for long-term storage because of gradual reagent oxidation).
- Select an optimal working concentration of the reagent and incubation time required for reagent deacetylation and oxidation for the specific cell line and assay conditions. If there are no protocols recommended for the specific cell line, start with a concentration from 1 to 10 μM and incubation for 30 min.
- Do not incubate the dye with the cells in the presence of serum because it contains enzymes that cleave H₂DCFDA.



外观:

分子 487.29

量:

CAS 4091-99-0

编号:

分子 C₂₄H₁₆Cl₂O₇

式:

IUPAC 2-(3,6-diacetyloxy-2,7-dichloro-9H-xanthen-9-yl)benzoic acid

名称:

溶解

度:

质量

控制:

储存条件: 本产品仅供研究目的提供和销售。 本产品并未经过食品、药品、医疗器械、化妆品等领域的安全性和效力测试，且未经明示或暗示授权用于其他任何用途，包括但不限于体外诊断、人类或动物用途，以及商业用途。

激发/吸收极大值，纳米: 511

ϵ , 摩尔吸光系数 $\text{m}^2\text{mol}^{-1}$: 118626

发射极大值，纳米: 533

荧光量子产率: 0.76

CF_{260} : 0.17

CF_{280} : 0.14