

Lumiprobe Corporation

201 International Circle, 135号套房 马里兰州亨特瓦雷, 21030

美国

电话: +1 888 973 6353 传真: +1 888 973 6354

电子邮件: order@lumiprobe.com

H2DCFDA (2',7'-dichlorodihydrofluorescein diacetate)

http://cn.lumiprobe.com/p/h2dcfda


 H_2DCFDA (2′,7′-dichlorodihydrofluorescein diacetate) is a common reagent used for investigating the production of reactive oxygen species in living cells.


 H_2DCFDA is a non-fluorescent fluorescein derivative (its reduced acetylated form). The reagent begins to emit fluorescence only after cleavage of acetyl groups and the reagent's oxidation in the cell while converting to 2',7'-dichlorofluorescein. This is a bright green-fluorescent dye (absorption maximum 511 nm, fluorescence maximum 533 nm). This reagent can be used for assays in living cells and is not compatible with sample fixation.

Acetyl groups in H_2DCFDA increase its lipophilicity and improve its cell membrane permeability. Once got into the cell, the dye is deacetylated by cell esterases, thus becoming charged and better fixed inside the cell. Oxidation with reactive oxygen forms results in the formation of a fluorescent product (2',7'-dichlorofluorescein) and can be detected using various methods, for example with a flow cytometer, plate reader, or fluorescent microscope.

Recommendations for using the reagent:

- Use a freshly prepared solution of the reagent (the working solution is not intended for longterm storage because of gradual reagent oxidation).
- Select an optimal working concentration of the reagent and incubation time required for reagent deacetylation and oxidation for the specific cell line and assay conditions. If there are no protocols recommended for the specific cell line, start with a concentration from 1 to 10 μ M and incubation for 30 min.
- Do not incubate the dye with the cells in the presence of serum because it contains enzymes that cleave H₂DCFDA.





外观:

分子量: 487.29 CAS 编号: 4091-99-0 分子式: $C_{24}H_{16}CI_2O_7$

IUPAC 名称: 2-(3,6-diacetyloxy-2,7-dichloro-9H-xanthen-9-yl)benzoic acid

溶解度: 质量控制: 储存条件: 激发/吸收极大值,纳米: 511 ϵ , 摩尔吸光系数, cm^{-1} : 118626

发射极大值,纳米: 533

荧光量子产率: 0.76

CF₂₆₀: 0.17

CF₂₈₀: 0.14