

(C18) Octadecanoyl-L-carnitine-d3

http://cn.lumiprobe.com/p/octadecanoyl-l-carnitine-d3

(C18) Octadecanoyl-L-carnitine-d3 chloride is a deuterated derivative of stearoyl-L-carnitine. The reagent is intended for use as an internal standard for the quantification of stearoyl-L-carnitine by GC- or LC-MS.

In addition, (C18) Octadecanoyl-L-carnitine d3 chloride, is suitable for the determination of (C18:1) Oleoyl-L-carnitine, (C18:2) Linoleoyl-L-carnitine, (C18:10H) 3-hydroxyoleoylcarnitine, (C18:20H) 3-hydroxylinoleoylcarnitine.

These acylcarnitines are biomarkers in the diagnosis of metabolic disorders such as carnitine palmitoyltransferase deficiency type I, type II, carnitine-acylcarnitine translocase (CACT) deficiency, Long-chain 3-hydroxyacyl-CoA dehydrogenase (LCHAD) deficiency, glutaric acidemia type II.



外 白色固体 观: 分 子 量: 467.14 CAS 2245711-27-5 编 号: 分 $\mathsf{C}_{25}\mathsf{H}_{47}\mathsf{D}_3\mathsf{CINO}_4$ 7子式: 溶 DMF,DMSO,乙醇 解 度: 质 NMR ¹H 和 HPLC-MS (95+%, D: 98+%) 量 里 控 制: 储 收到后 -20°C 避光保存 24 个月。运输: 室温下最多可保存3周。干燥。 晶存条件: 法律声 本产品仅供研究目的提供和销售。 本产品并未经过食品、药品、医疗器械、化妆品等领域的安全性和效力测试,且未经明示或暗示授权用于其他任何用途,包括但不限于体外诊 断、人类或动物用途,以及商业用途。 , 明: