

3-(5-methyl-1H-pyrazol-3-yl)propionic acid-d6 (MPP)


<http://cn.lumiprobe.com/p/sa-succinylacetone-pyrazol-d6>

Succinylacetone (abbreviated as SA) is a product of tyrosine catabolism, and an inhibitor of heme biosynthesis. Succinylacetone testing in dried neonatal blood spots, followed by quantitation of SA in blood or urine in high-risk neonates, has excellent sensitivity and specificity for the diagnosis of tyrosinemia type 1 [1].

Succinylacetone is a reactive diketone which reacts with the amino groups of peptides and proteins present in the blood. Therefore, extraction of the bound SA requires its conversion into a product suitable for its extraction and quantification. This is effected by means of hydrazine treatment which leads to a conversion of succinylacetone to this pyrazole derivative.

This pyrazole derivative is a deuterated derivative is used as an analytical standard in the measurement of succinylacetone (SA) in dried blood spot samples.

[1] Kehar M., Sen Sarma M., Seetharaman J., Jimenez Rivera C., Chakraborty P. Decoding hepatorenal tyrosinemia type 1: Unraveling the impact of early detection, NTBC, and the role of liver transplantation. Can Liver J. 2024. 7(1). P.54-63.



外观 白色固体

分子量 160.21

分子式 $C_7H_4D_6N_2O_2$

溶解度 在水中

质量控制 收到后 -20°C 避光保存 24 个月。运输：室温下最多可保存3周。干燥。

储存条件

法律声明 本产品仅供研究目的提供和销售。本产品并未经过食品、药品、医疗器械、化妆品等领域的安全性和效力测试，且未经明示或暗示授权用于其他任何用途，包括但不限于体外诊断、人类或动物用途，以及商业用途。