



## ssGreen RNA Gel Staining Solution, 10,000×

<http://cn.lumiprobe.com/p/ssgreen-ii-gel-stain>

ssGreen® is one of the most sensitive stains for post-electrophoresis staining of RNA and single-stranded DNA (ssDNA) in agarose or polyacrylamide gels. The fluorescence quantum yield of the ssGreen/RNA complex is more than 7× higher than that of the ethidium bromide/RNA complex. Although ssGreen is not selective for RNA staining, the dye exhibits about 1.5× greater quantum yield when bound to RNA than double-stranded DNA, which makes it unique among all nucleic acid dyes.

ssGreen staining is compatible with denaturing gels. On agarose/formaldehyde and polyacrylamide/urea gels, the sensitivity of ssGreen is slightly reduced but still superior to that of ethidium bromide. Staining agarose/formaldehyde gels with ssGreen does not interfere with transfer of RNA to filters or subsequent hybridization in Northern blot analysis as long as 0.1%-0.3% SDS is included in prehybridization and hybridization buffers.

Staining gels with ssGreen has fewer steps than those with ethidium bromide. Because the fluorescence of ssGreen/RNA complexes is not quenched by formaldehyde or urea, there is no need to wash these denaturants out of gels before staining. Also, ssGreen stain has a low intrinsic fluorescence, allowing gel viewing and photographing without preliminary removing unbound dye.



外观: 深橙色溶液

分子量: 581.52

CAS 编号: 172827-25-7

分子式: C<sub>28</sub>H<sub>28</sub>N<sub>3</sub>OS

质量: 分光光度法

控制:

储存: 收到后在 -20°C 黑暗条件下可保存 24 个月。运输: 室温下最多可保存 3 周。干燥。

条件:

说明:

法律: 本产品仅供研究目的提供和销售。本产品并未经过食品、药品、医疗器械、化妆品等领域的安全性和效力测试, 且未经明示或暗示授权用于其他任何用途, 包括但不限于体外诊断、人类或动物用途, 以及商业用途。

声明:

激发/

吸收

极大

值:

纳

米:

483 (complex)

发射 518 (complex)  
极大  
值,  
纳  
米: